

This Page Is Inserted by IFW Operations  
and is not a part of the Official Record

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problem Mailbox.**

Gram Staining of isolate TW-S-7-1






FIG. 1



Electrophoresis of avian collagen on a 7.5% polyacrylamide gel.  $\alpha$  and  $\beta$  denote different collagen configurations.


**FIG. 2**



Separation of porcine collagen on a 10% polyacrylamide gel.  $\alpha$ ,  $\beta$  and  $\gamma$  denote different collagen configurations.

Lanes 1, 2 porcine collagen, 20  $\mu$ g and 40  $\mu$ g; lanes 3, 4 commercial bovine collagen, 40  $\mu$ g and 20  $\mu$ g

**FIG. 3**



8% SDS-PAGE


Lane 1:

10  $\mu$ l

Lane 2:

20  $\mu$ l

**FIG. 4**



8% SDS-PAGE


Lane 1: 5  $\mu$ l

Lane 2: 10  $\mu$ l

Lane 3: 20  $\mu$ l

Lane 4: 40  $\mu$ l

**FIG. 5**



8% PAGE-SDS

Lane 1 : marker

Lane 2 : 5  $\mu$ l

Lane 3 : 10  $\mu$ l

**FIG. 6**